

Oxidative Stress in Diabetes and Periodontitis

Janaina de Cássia Orlandi Sardi

Department of Clinical Analysis, Laboratory of Clinical Mycology, Faculty of Pharmaceutical Sciences,
Univ Estadual Paulista, Araraquara, São Paulo, Brazil

Address for correspondence: Dr. Janaina de Cássia Orlandi Sardi, Department of Clinical Analysis, Laboratory of Clinical Mycology, Faculty of Pharmaceutical Sciences, Univ Estadual Paulista, Araraquara, São Paulo, Brazil. E-mail: janaina-sardi@uol.com.br

Oxidative stress is an imbalance between the production of a reactive oxygen species and the antioxidant defense, leading to tissue damage. The produced reactive oxygen species, such as superoxide anion, hydroxyl radical, and peroxy radical result in damage to many biological molecules (including DNA, lipids, and protein), and the prolonged existence of these reactive oxygen species promotes severe tissue damage and cell death.^[1,2] It has been proposed that there is a causal relationship between insulin resistance, oxidative stress, and periodontitis and that hyperglycemia is a major factor responsible for the activation of oxidative stress.^[1,2]

The presence of malondialdehyde, the products of the hydroxylation of DNA bases such as 8-hydroxy-2'-deoxyguanosine (8-OHdG) were found in crevicular fluid and periodontal pockets.^[3] Cell damage is a consequence of the effect of free radicals and chronic oxidative stress.^[3] Furthermore, Sawamoto *et al.*,^[4] proposed a close relationship between 8-OHdG and periodontal pathogens. When further considering periodontal disease, hypoxia of the tissue occurs at the onset of periodontitis,^[5] which stimulates the production of cytokines and inflammatory mediators involved in alveolar bone resorption, including IL-6 and IL-1. With re-oxygenation after hypoxic events, there is an increased level of superoxide anions and other reactive oxygen species, which in turn induce degradation of the NF- κ B inhibitor, resulting in greater tissue destruction. Therefore, periodontal disease associated with diabetes,

smoking, and/or occlusal trauma leads to periodontal tissue hypoxia and re-oxygenation events, a situation that hypothetically increases the morbidity of periodontal pathology.

Neutrophils in poorly controlled diabetic patients released significantly more superoxide than neutrophils from patients with good glycemic control and from non-diabetic healthy individuals.^[6] Neutrophils increased protein kinase C activity, elevated amounts of diacylglycerol, and enhanced nicotinamide adenine dinucleotide phosphate oxidase activity. This suggests that hyperglycemia can lead to neutrophil activation and elevated protein kinase C activity, resulting in increased oxidative stress.^[6]

Pendyala *et al.*,^[7] demonstrated that increasing oxidative stress can be an important contributing factor in the pathogenesis of diabetes and periodontal disease. Therefore, the co-existence of these conditions could pathologically increase the effect of oxidative stress.

References

1. Nassar H, Kantarci A, van Dyke TE. Diabetic periodontitis: A model for activated innate immunity and impaired resolution of inflammation. *Periodontol* 2000 2007;43:233-44.
2. Mahadev K, Wu X, Zilbering A, Zhu L, Lawrence JT, Goldstein BJ. Hydrogen peroxide generated during cellular insulin stimulation is integral to activation of the distal insulin signaling cascade in 3T3-L1 adipocytes. *J Biol Chem* 2001;276:48662-9.
3. Marfella R, Quagliaro L, Nappo F, Ceriello A, Giugliano D. Acute hyperglycemia induces an oxidative stress in healthy subjects. *J Clin Invest* 2001;108:635-6.
4. Sawamoto Y, Sugano N, Tanaka H, Ito K. Detection of periodontopathic bacteria and an oxidative stress marker in saliva from periodontitis patients. *Oral Microbiol Immunol* 2005;20:216-20.
5. Motohira H, Hayashi J, Tatsumi J, Tajima M, Sakagami H, Shin K. Hypoxia and reoxygenation augment bone-resorbing

Access this article online

Quick Response Code:



Website:
www.najms.org

- factor production from human periodontal ligament cells. J Periodontol 2007;78:1803-9.
6. Karima M, Kantarci A, Ohira T, Hasturk H, Jones VL, Nam B-H, *et al.* Enhanced superoxide release and elevated protein Kinase C activity in neutrophil from diabetic patients: Association with periodontitis. J Leukocyte Biol 2005;78:1-9.
 7. Pendyala G, Thomas B, Joshi SR. Evaluation of Total

Antioxidant Capacity of Saliva in Type 2 Diabetic Patients with and without Periodontal Disease: A Case-Control Study. N Am J Med Sci 2013;5:51-7.

How to cite this article: Sardi JO. Oxidative stress in diabetes and periodontitis. North Am J Med Sci 2013;5:58-9.

Source of Support: Nil. **Conflict of Interest:** None declared.

New features on the journal's website

Optimized content for mobile and hand-held devices

HTML pages have been optimized of mobile and other hand-held devices (such as iPad, Kindle, iPod) for faster browsing speed.

Click on **[Mobile Full text]** from Table of Contents page.

This is simple HTML version for faster download on mobiles (if viewed on desktop, it will be automatically redirected to full HTML version)

E-Pub for hand-held devices

EPUB is an open e-book standard recommended by The International Digital Publishing Forum which is designed for reflowable content i.e. the text display can be optimized for a particular display device.

Click on **[EPub]** from Table of Contents page.

There are various e-Pub readers such as for Windows: Digital Editions, OS X: Calibre/Bookworm, iPhone/iPod Touch/iPad: Stanza, and Linux: Calibre/Bookworm.

E-Book for desktop

One can also see the entire issue as printed here in a 'flip book' version on desktops.

Links are available from Current Issue as well as Archives pages.

Click on  View as eBook